


Journal Pre-proof

Effect of Adding Global Postural Re-education to Kendall Exercises for Treating Asymptomatic Forward Head Posture: A Single-Blinded Randomized Controlled Trial

Walaa Abu-Taleb, Abeer Abdelrahman Yamany, Yasser M. Aneis, Shimaa T. Abu El Kasem



PII: S1360-8592(25)00039-7

DOI: <https://doi.org/10.1016/j.jbmt.2025.01.039>

Reference: YJBMT 3296

To appear in: *Journal of Bodywork & Movement Therapies*

Received Date: 11 August 2024

Revised Date: 2 January 2025

Accepted Date: 22 January 2025

Please cite this article as: Abu-Taleb, W., Yamany, A.A., Aneis, Y.M., Abu El Kasem, S.T., Effect of Adding Global Postural Re-education to Kendall Exercises for Treating Asymptomatic Forward Head Posture: A Single-Blinded Randomized Controlled Trial, *Journal of Bodywork & Movement Therapies*, <https://doi.org/10.1016/j.jbmt.2025.01.039>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2025 The Author(s). Published by Elsevier Ltd.

Title

Effect of Adding Global Postural Re-education to Kendall Exercises for Treating Asymptomatic Forward Head Posture: A Single-Blinded Randomized Controlled Trial.

Authors

1. Walaa Abu-Taleb a. Physiotherapy Department, School of Human Sciences, London Metropolitan University, London, United Kingdom and b. Department of Basic Science, Faculty of Physical Therapy, Cairo University, Giza, Egypt. Email: w.abutaleb11@gmail.com
2. Abeer Abdelrahman Yamany, b. Department of Basic Science, Faculty of Physical Therapy, Cairo University, Giza, Egypt. Email: abeer.yamany@pt.cu.edu.eg
3. Yasser M. Aneis, b. Department of Basic Science, Faculty of Physical Therapy, Cairo University, Giza, Egypt. Email: dryassercom@yahoo.com
4. Shimaa T. Abu El Kasem, b. Department of Basic Science, Faculty of Physical Therapy, Cairo University, Giza, Egypt and c. Department of Basic Science, Faculty of Physical Therapy, Galala University, Suez, Egypt. Email: shaimaataha@cu.edu.eg

Corresponding Author

Walaa Abu-Taleb
School of Human Sciences
London Metropolitan University
London, United Kingdom
w.abutaleb11@gmail.com

Effect of Adding Global Postural Re-education to Kendall Exercises for Treating Asymptomatic Forward Head Posture: A Single-Blinded Randomized Controlled Trial

ABSTRACT

Introduction Forward head posture (FHP) is a common postural malalignment in young population that is associated with limitation of mobility and functional disability. Kendall exercises are one of the commonly used postural correction techniques to treat FHP. Global Postural reeducation (GPR) is a postural correction exercise commonly used for musculoskeletal disorders. The current study aimed to investigate the combined effect of GPR and Kendall Exercises in the treatment of FHP.

Methods A single-blinded parallel-groups randomized controlled trial was conducted. Forty-three subjects aged 18-30 years were recruited with FHP marked by a craniovertebral angle (CVA) less than 50°. Participants were randomly allocated into two groups: group A (GrA) received GPR plus Kendall Exercises, and group B (GrB) received Kendall Exercises only. Variables were measured before and immediately after 12 sessions of treatment including CVA, gaze angle (GA), shoulder angle (SA), cervical range of motion (CROM), neck disability index (NDI), chest expansion, and spinal mobility.

Results Between groups analysis revealed no statistically significant difference between either treatment in CVA, CROM, and NDI. There was a statistically significant improvement of chest expansion and spinal mobility in favor to GrA. Within-group analysis revealed that both interventions were statistically significant in improving CVA, CROM, and NDI ($P < 0.05$). Both treatments showed no statistical difference in GA and SA.

Conclusions The added effect GPR technique to Kendall exercises significantly improved CVA, CROM, NDI, chest expansion, and spinal mobility in subjects with FHP.

Keywords Forward Head Posture, Global Postural Re-education, Kendall Exercise, Postural Correction, Craniovertebral Angle.

INTRODUCTION

Forward head posture (FHP) is a common non-structural malalignment of the cervical spine characterized by hyperflexion of the lower cervical spine (C2-C7), and hyperextension of the upper cervical vertebrae which includes the occiput, atlas and axis (C0-C2) (Khayatzaheh et al., 2017). FHP is notably spread among young adults and university students with a prevalence of 63.3% to 90% in the population (Karthik et al., 2022; Kose et al., 2022). Some of the underlying causes of FHP include faulty postural habits, sedentary lifestyle and prolonged use of smart devices (Fercho et al., 2023).

The anterior head translation in FHP causes compression of the cervical vertebrae (Bonney & Corlett, 2002), which triggers overactivity of the cervical and suboccipital muscles that is required to counterbalance the weight of the head (Alowa & Elsayed, 2021). The muscles overactivity limits the cervical mobility and further increases the load on the vertebrae and intervertebral discs which raises the risk of spinal injury (Alowa & Elsayed, 2021; Bonney & Corlett, 2002; Sarraf & Varmazyar, 2022). The disturbance of the normal cervical mechanics extends to the thorax, resulting in diaphragmatic and respiratory dysfunction (Koseki et al., 2019). FHP is

commonly associated with restricted cervical and chest mobility (Kim et al., 2018), impaired cervical neuromotor control (Khan et al., 2020), and declined pulmonary function (Koseki et al., 2019).

Correcting FHP and related symptoms are attainable by various corrective exercises and manual therapy techniques (Fathollahnejad et al., 2019; Sheikhhoseini et al., 2018). Kendall Exercises (KE) were first described by Florance Kendall to correct the acquired postural fault of FHP, kyphosis, and rounded shoulders. This technique is comprised of four exercises including: stretching the tight neck extensors and pectoral muscles and strengthening the weak deep neck flexors and scapular retractors (Kendall et al., 2005). Kendall's technique is recommended to correct head and shoulders faulty postures and improve mobility and functional ability of the neck and shoulder complex (Harman et al., 2005; Kim et al., 2015; Kong et al., 2017; Rahul S et al., 2024).

Global Postural Re-education (GPR) is another spinal postural correction technique that uses a series of active symmetrical lengthening positions maintained by antagonists muscles contraction with breathing training (Pillastrini et al., 2018; Souchard et al., 2011). The postures of GPR restore a balanced activation between antero-posterior postural muscles

within myofascial chains while avoiding substitutions (Mendes-Fernandes et al., 2021; Souchard et al., 2011). Maintaining the stretching position for 15-20 minutes in GPR is necessary to engage the viscoelastic trait “creep” of the tissues, to stimulate a reflex relaxation of the agonists and to increase the stretch tolerance (Fukaya et al., 2022; Souchard et al., 2011). GPR has shown significant effectiveness in the treatment of postural impairments and musculoskeletal (MSK) conditions with reported value in decreasing pain, restoring mobility, and decreasing physical and pulmonary functional disabilities (Carrasco-Lopez & Medina-Porqueres, 2016; Dimitrova & Rohleva, 2014; Ferreira et al., 2016).

Three building principles support the GPR technique: Individuality (Individualité), where a personalized care is offered to patients, Causality (Causalité), where a MSK condition may be caused by a distant area, and Globality (Globalité), where the body is connected via myofascial chains and must be treated wholistically (Ferreira et al., 2016; Souchard et al., 2011). The globality principle in GPR may be supported by the contemporary research around fascia (Wilke et al., 2016; Wilke & Krause, 2019). The fascial system extends across the whole body to support and connect the different elements of the MSK system, creating an interconnected web of myofascial chains (Blottner et al., 2019; Bordoni &

Myers, 2020). The fascial continuum can retain and transmit up to 50% of the muscles' tension, which allows the fascia to coordinate the interplay of movements across the myofascial chains longitudinally and transversely in local and global areas via what is known as myofascial force transmission (Bordoni & Myers, 2020; Findley et al., 2015; Huijing, 2009; Stecco et al., 2023; Wilke et al., 2018).

Empirical research has investigated the applicability of myofascial force transmission. In the study by Wilke et al. (2017) stretching exercise applied proximally (cervical) or distally (calf) was equally effective in improving cervical mobility and supporting the value using myofascial force transmission in rehabilitation. Several studies further compared the superiority of global techniques (e.g. GPR) and local techniques in the rehabilitation of various MSK conditions (De Amorim et al., 2014; França et al., 2012; Matsutani et al., 2023; Mendes Fernandes et al., 2023; Rosário et al., 2012). However, the studies did not lead to a consensus on the comparability of these techniques or the value of combining both treatments in MSK conditions. The proven benefits of using Kendall Exercises added to the hypothesized efficacy of GPR may lead to better outcomes in patients with FHP. To the best of the authors' knowledge, no study considered adding GPR to local treatment as a treatment approach in the treatment of

FHP. This study aimed at assessing the effect of GPR added to KE on postural angles (craniovertebral angle (CVA), gaze angle (GA), and shoulder angle (SA)), cervical range of motion (CROM), neck disability index (NDI), chest expansion (CE), and spinal mobility in people with FHP.

MATERIALS AND METHODS

Design

The current study was designed as a prospective single-blinded parallel-group RCT that was conducted between September 2020 - October 2021. The study followed the recommendations of CONSORT for randomised clinical trials (Figure 1). The study was approved and registered by the research ethical committee of the Faculty of Physical Therapy, Cairo University, Giza, Egypt (P.T.REC/012/002671) and registered in ClinicalTrials.gov (NCT04723511).

Subjects

Subjects were recruited via electronic and written adverts in the faculty of physiotherapy at Horus University in Egypt with the opportunity to be screened for FHP for a research project where those identified with FHP were invited to participate in this study. Subjects were recruited from both sexes, age 18-30 years old, asymptomatic or neck pain <3 on the visual

analogue scale (VAS) (the score below 4 is interpreted as no pain) (Jensen et al., 2003), CVA < 50 degrees (Moustafa et al., 2020), and body mass index (BMI) ranges from 18.5-28. Participants were excluded from the trial if they had a history of traumatic neck injury, scoliosis, severe temporomandibular or visual problem, spinal surgeries or fixations, severe respiratory condition, daily screen time for >4 hours (Jung et al., 2016), a neurologic deficit (Cunha et al., 2008), or received any treatment for the neck in the last three months.

Sample size

Sample size calculation was performed using G*Power 3.0.20 software (Faul et al., 2007), using the results of the primary outcome CVA from a pilot study. The pilot study was conducted on 10 subjects with FHP who received the same interventions and divided equally between the experimental and control groups. T-test was used within and between interaction, the effect size = 0.8, $\beta = 0.2$, and $\alpha = 0.05$, revealed that 38 (19 per group) subjects were the appropriate sample size. Five subjects were then added to compensate for any withdrawals. The appropriate sample was 43 subjects divided into 2 groups.

Randomization

Seventy-eight subjects were recruited and screened for FHP using measurement of CVA via photogrammetry and kinovea. The subjects who

did not fulfill the inclusion criteria (n=35) were excluded; the details of this number can be followed up in Figure 1. The randomization was carried out using a computer-generated numbers in opaque sealed envelopes. Subjects were assigned to one of the two groups in a 1:1 ratio. Allocation was performed by one of the authors who was not involved in either data collection or treatment. To maintain allocation concealment, envelopes were opened only at the time of enrolment of each subject. The subjects were blinded to their allocation to minimize bias until the study concluded (Page & Persch, 2013). The procedure was explained and informed consent forms were then signed by the participants.

Interventions

The intervention was provided for 12 treatment sessions over four weeks (Im et al., 2016), by a physiotherapist who has over 15 years of MSK clinical experience. As a general rule, subjects were instructed to work ergonomically and practice treatment exercises at home, and to avoid poor posture and excessive smart devices exposure. An ergonomic online training was provided by the principal researcher and the exercises were printed for all the subjects.

Kendall Exercises were performed actively by all the subjects (completed in 20-30 minutes) while the therapist observed the execution of the exercises, corrected faulty performance and timed the stretching exercises with a stopwatch. The difficulty of the exercises was gradually incremented if the subject can complete 10 correct exercise repetitions for three sets with ease (Harman et al., 2005; Kim et al., 2015). The description, dose, and progression of KE is available in Table 1 (Kendall et al., 2005; Kim et al., 2015).

GPR sessions started with five minutes of gentle diaphragmatic release and training of deep diaphragmatic breathing to stretch the diaphragmatic (respiratory) chain. Diaphragmatic release involved applying gentle pressure with the fingertips of the therapist starting from the xyphoid process moving towards the lateral lower ribs and was repeated three times (Moreno, Catai, Teodori, Borges, De Cesar, et al., 2007). Diaphragmatic breathing was performed actively in supine and the therapist applied cervical traction and corrected accessory muscles activation. It was performed five times at the start of the session then repeated three times every five minutes during the GPR positions.

The GPR postures selected in this trial aimed to stretch the anterior and posterior chains commonly affected in FHP (Fernández-De-Las-Peñas

et al., 2005). Postures difficulty (lying to standing) and the amount of tension applied were incrementally increased as the treatment progressed to allow the subject to gain more strength and motor control and to build stretch tolerance (Mendes-Fernandes et al., 2021; Souchard et al., 2011). During the treatment, a cephalic traction of the head and caudal traction at sacral levels were passively applied by the therapist to flatten and elongate the spine, which was repeated during the session as the therapist deemed necessary (Silva et al., 2012). Treatment time of GPR ranged from 35-70 minutes according to the progression phase. Postures images are displayed in Figure 2 while details of the GPR technique applied are in Table 2.

To enhance the therapist's ability to provide accurate, personalized, and symmetrical postural correction in GPR positions, a smartphone application with gridline view was used. ACPP Core2 (Jinnyu Technology CO, LTD), is free smartphone application that displays a calibrated gridline with squares of 0.25 cm^2 area and a plump line to allow for posture symmetry that was not detectable by the naked eye. The subjects were instructed to assume the GPR postures as symmetrically as possible, and a photo was taken. The researcher then used the ACPP Core2 to correct the postures and take a photo of the corrected position. The photos were then shown to the subjects to reinforce visual feedback of correct the posture (Xu et al., 2019). Body awareness training is an effective non-invasive treatment in postural

deviations that could improve posture anomaly, postural control and vertical perception (Yagci et al., 2018).

Outcome Measures

Evaluations were performed before and after the completion of the treatment sessions. The assessments were performed by one of the authors that was not involved in randomization or treatment process. The primary outcome measure for this study was Craniovertebral Angle (CVA) while the secondary outcome measures included gaze angle (GA) and shoulder angle (SA) measured by photogrammetry and analyzed by Kinovea software, cervical range of motion (CROM) measured by cervical range of motion instrument, neck function measured by neck disability index (NDI), and a tape measure was used to measure both the circumference of the chest expansion (CE) and the spinal mobility via fingers to floor test (FTF).

postural angles and photogrammetry

Photogrammetry is a valid and reliable method to measure postural angles (Singla et al., 2017). Three angles were assessed in this study including CVA for forward head posture, GA for upper cervical posture, and SA for rounded shoulders (See Figure 3) (Singla et al., 2017). The assessed postural angles were considered normal when $CVA > 50^\circ$, $GA < 15^\circ$ and SA

> 52° (Moustafa et al., 2020; Ruivo et al., 2017; Van Det et al., 2008). The photographs were taken with a smartphone's Camera (12-megapixel) mounted on the tripod 1.5 meters away from the subjects and the height was adjusted to each subject's neck (Puig-Diví et al., 2019). Self-adhesive markers were placed on C7 and the acromion processes bilaterally to improve the visualization of bony landmarks in the captured photos. Six photos from the sagittal planes were taken for each subject in standing position. Analysis of photogrammetry was performed via open-access Kinovea Software (Version 0.9.5) under the GPL v2 license (Charmant & Contributor, 2021). Kinovea is valid, precise, and reliable for angles and distances (Lee et al., 2017; Puig-Diví et al., 2019). The measurement technique used followed the guidance of Mun Cheung Lau et al. (2010).

cervical mobility

Cervical range of motion instrument has good validity and reliability in the assessment of neck mobility 95% confidence interval (CI), ICC= 0.89-0.98 (Audette et al., 2010; Tousignant et al., 2006). Active CROM was measured in flexion, extension, bilateral side bending, and bilateral rotation. Following familiarization with the procedures and warming up, CROM was assessed according to the procedures explained by Wang et al. (2002). The measurements were repeated 3 times, and the average was calculated. Any

contaminated trial by trunk or shoulder substitution was discarded and replaced with another (Wang et al., 2002).

neck function

(Macdelilid et al., 2009) reported that NDI is a valid and reliable scale that can be used for acute and chronic MSK dysfunctional conditions. NDI for mechanical neck pain without upper extremity symptoms exhibited excellent reliability 95% CI, ICC = 0.88 (Pt et al., 2018). NDI is a self-administered test for the neck functional assessment in 10 activities of daily living (Ackelman & Lindgren, 2002). Each of the activities is scored by progressive statements from no disability to maximum disability. Each statement is then scored from zero to 5, where the zero equates not having any disability and 5 stands for maximum disability (Vernon & Mior, 1991). Subjects in this trial were instructed to select the statement that represents them best.

chest expansion

Chest Expansion/mobility (CE) via tape measure has shown to be valid and highly reliable for the upper and lower chest expansion indicating chest mobility and indirectly lung function (Najwatul et al., 2016). Tape measure has a very good interrater and intratester reliability measuring upper and lower chest expansion with 95% CI, ICC= 0.78- 0.84, and 0.85-0.93 respectively (Reddy et al., 2019). Assessment of the chest expansion was

done at two levels: upper thoracic or axial at the level of the armpit, and lower thoracic/xiphoid at the level of the xiphoid process (Bennett et al., 2021; Mohan et al., 2012; Reddy et al., 2019). The measurement procedures were done following the approach of Reddy et al. (2019) and the average of three trials were calculated.

spinal mobility

Fingertips to floor test (FTF) was used to assess the spinal mobility. FTF test is a valid and reliable test with a 95% CI, ICC= 0.96 (Hecimovich & Hebert, 2016; Perret et al., 2001), and has a strong correlation to spinal mobility (Guo et al., 2023). The test involves measuring the distance between the middle finger to the floor without knee flexion via tape measure (Perret et al., 2001).

Statistical Analysis

An unpaired t-test was conducted for the comparison of subjects' demographic characteristics between groups and Chi-squared test was used for comparison of sex distribution. Shapiro-Wilk test was used to test normality of the data while the homogeneity of variances between groups was confirmed by Levene's test. Mixed MANOVA was performed to compare within and between groups effects of the treatment interventions on CVA, GA, SA, CROM, CE (axillary and xiphoid) and spinal mobility (FTF

test). Bonferroni corrections were carried out for subsequent multiple comparisons. Mann–Whitney test was used for comparison of NDI between groups and Wilcoxon signed ranks test was used for comparison between pre- and post-treatment in each group. The statistical analysis was completed using the statistical package for social studies (SPSS) version 25 for Windows (IBM SPSS, Chicago, IL, USA). The level of significance for all statistical tests was set at $p < 0.05$.

RESULTS

Forty-three subjects completed the study without drop-out, hazards or adverse effects reported from the subjects during treatment. Table (3) shows demographic characteristics of both GrA and GrB. There was a match at the baseline between groups in age, sex and BMI $p > 0.05$. Mixed MANOVA revealed that there was a significant interaction between treatment*time ($F=13.38$, $p=0.001$). There was also a significant main effect of time ($F=23.45$, $p=0.001$), but there was no significant main effect of treatment ($F=0.99$, $p=0.47$).

Between group analysis revealed that there was no significant difference between groups pretreatment ($p > 0.05$). There was no statistically significant difference in CVA, GA, SA, CROM and NDI between groups post-treatment ($p > 0.05$). There was a significant increase in axillary and

sternal circumferences, and a significant decrease in FTF of GrA compared with GrB ($p < 0.01$) (Table 4 and 5).

Within group analysis revealed that there was a post-treatment significant statistical improvement in both groups in CVA, CROM ($p < 0.01$), and NDI ($p < 0.05$) compared with pretreatment. There was a significant increase in axillary and sternal CE and a significant decrease in FTF in GrA ($p < 0.001$), while there was no significant change in GrB ($p > 0.05$). There was no significant difference in post treatment measurements of GA and SA in both groups ($p > 0.05$). Table 5 shows the significant decrease in NDI of groups A and B post-treatment compared with pretreatment.

DISCUSSION

The current study aimed to identify the effect of adding GPR to KE on postural angles (CVA, GA and SA), cervical mobility (CROM) and function (NDI), chest expansion (CE), and spinal mobility (FTF) in patients with FHP. The results showed a statistically significant difference in favor to the experimental group (GrA) only in CE and FTF. Within groups analysis showed significant statistical improvement in both groups in CVA, CROM, and NDI while there was not any significant difference in GA and SA.

The measures of CE were improved by 5.69 cm and 7.04 cm in axial and xyphoid diameters respectively. The improvement exceeded the minimum clinically important difference (MCID) of 3.60 cm for the axial CE and 4.40 cm for the xyphoid CE (Reddy et al., 2019). This improvement may be facilitated by the adoption of GPR positions and repeated breathing workout which corrects the mechanical fault of the neck and thorax and restores the normal function of respiratory muscles and diaphragm (Koseki et al., 2019; Rocha et al., 2018). The breathing workout in GPR is postulated to increase the recruitment of respiratory muscle fibers while simultaneously stretching the diaphragmatic chain and augmenting the extensibility of the diaphragm (Rocha et al., 2018). In agreement with the current study, GPR was reported as beneficial in augmenting the respiratory muscle strength, chest mobility, maximal respiratory pressure and reducing pain among other benefits in the rehabilitation of MSK dysfunction (Moreno, Catai, Teodori, Borges, De Castro Cesar, et al., 2007).

FTF test is typically utilized in clinical setting and in research as a screening tool for spinal mobility in low back pain and ankylosing spondylitis patients (Bonetti et al., 2010; Silva et al., 2012). However, it was used in the current study, similar to that by Cavalcanti et al. (2020) as an indicator of the globalized effect of GPR stretching exercises on posterior myofascial chain that extends from upper trapezius to the foot intrinsic

muscles (Bonetti et al., 2010). Our results showed a significant improvement of FTF scores from baseline by 7.68 cm which exceeded the minimal detectable change of 4.5 cm (Guo et al., 2023). The improvement of FTF following GPR could be a result of stretching the posterior myofascial chain for a prolonged time using GPR positions (Bonetti et al., 2010). This is in agreement with the previous results of Cavalcanti et al. (2020) who found similar decline of FTF scores following 10 sessions of GPR in their pilot study of 18 subjects with spinal pain despite not showing any change in posture.

Further, GPR uses isometric contraction of the antagonistic muscles which may cause a reflex inhibition of agonists and facilitate increased the length of muscles (Silva et al., 2012). This phenomenon is known as reciprocal inhibition which is commonly used to explain the value of proprioceptive neuromuscular facilitation stretching technique (Hindle et al., 2012). Furthermore, Oliveri et al. (2012) investigated the effect of GPR standing position and reported an inhibition of the motor cortical area controlling the agonists muscles and an activation of cortical regions controlling the antagonists which further supports the proposed effect of GPR.

Our results showed no improvement of GA and SA as a result of GPR and KE treatments despite improving CVA which indicates the correction of FHP. This contrasts the findings of previous studies that utilized KE to correct FHP (Heydari et al., 2022; Kim et al., 2015). This study assumes that as the inclusion criteria of the participants were only marked by CVA angle not GA or SA, the awareness of the participants' body behavior, and postural habits which in turn affects self-education and correction may have not extended to the upper cervical or shoulder (Schwertner et al., 2018; Siemonsma et al., 2013). Another possible explanation of the lack of improvement in GA might be provided in the findings of Lin et al. (2022) who studied cadavers with mild FHP and found that rectus capitis (occipital DNF) was in a shortened position while longus capitis (cervical DNF) was in a lengthened position. This may suggest that applying exercises to activate both of these muscles equally might not evidently be the best option to improve cervical and head position (CVA and GA) simultaneously (Lin et al., 2022).

The results of this study showed significant improvement of CVA, CROM and NDI in both groups. The post-treatment CVA scores exceeded the cutoff value of 50° that identifies subjects with FHP in both groups by ($\geq 6^\circ$, $p=0.001$), and surpassed the MCID in photogrammetry at 1.40° (Heydari et al., 2022; Lee et al., 2016). Further, CROM also improved in

both groups by (5.33° - 8.75° , $p = 0.001$) which lies within the reported MCID 3° - 10 (Jørgensen et al., 2017; Williams et al., 2010). A significant improvement of NDI after treatment is evident in both groups ($p = 0.001$).

The improvements of CVA, CROM, and NDI in the current study may be attributed to the effect of normalizing the cervical curve. Two FHP correction techniques were used in this study to re-establish the normal antero-posterior muscle balance: Kendall technique work by strengthening the weak deep neck flexors (DNF) and scapular retractors and stretching the tight neck extensors and shoulder protractors. While GPR technique applies active spinal lengthening technique with breathing control that is further accentuated by a passive manual cervical/ sacral traction by the therapist. Both techniques encompass DNF activation, that is renowned for normalizing the cervical curve in FHP (Alowa & Elsayed, 2021; Chang et al., 2023).

Activating DNF, particularly longus capitis and longus colli, was found in functional MRI to cause flexion of the cranio-cervical junction and flattening of cervical lordosis (Cagnie et al., 2008). Further, strong DNF allows relaxation of the overly active accessory muscle that contribute to the maintenance of FHP and fosters restoration of cervical mobility and

functionality (Alowa & Elsayed, 2021; Kong et al., 2017). The normalization of the curve would then correct the biomechanical faults of the cervical spine and leads to improvement of limited ROM and cervical dysfunction.

Previous studies found that smaller CVA is associated to with higher NDI score and lower CROM (De-La-Llave-Rincón et al., 2009; Kinjal Bagthariya & Kakkad, 2024; Yip et al., 2008). Similarly, Chang et al. (2023) reported in their systematic review that correcting the postural deviation of FHP and rounded shoulders is reflected in restoring cervical mobility and improving both pain and functional disability (Chang et al., 2023). However, it is important to remember that these interpretations are correlational not causal.

As evident in our findings are comparable to previous studies that reported and improvement of CVA, cervical mobility, and functional disability as a result of utilizing the Kendall technique (K.-H. Kim et al., 2015; Kong et al., 2017). On the other hand, no previous studies were found assessing the use of GPR for FHP but it was repeatedly used for cervical pain other spinal conditions. GPR is found to be fruitful in improving VAS, NDI, CROM and neck muscle electrical activity (Cunha et al., 2008; De Amorim et al., 2014; Mendes Fernandes et al., 2023; Pillastrini et al., 2018).

This randomized controlled study addressed an important issue related to the effectiveness of combined global and local treatment for asymptomatic FHP. The strengths of this study lies in its design, diverse sample, limitation of confounding factors, randomization, and the use of valid and reliable measurement tools all increase the robustness of our results and decreases bias in our findings.

Future research needs to address the limitations in this study that may limit the generalizability of the findings. The short-term (<four weeks) or the long-term effects (>four weeks) of the combined treatment techniques on cervical spine were not studied. Also, the effect of treatment on the thoracic or lumbar spines was not investigated. Furthermore, we did not investigate patient satisfaction and the cost and time effectiveness of both techniques in relation to achieved results.

CONCLUSION

The combined effect of GPR and local KE for 12 sessions has showed a significant improvement of FHP angle, cervical mobility, neck functionality, axial and xyphoid chest expansion, and spinal mobility in patients with FHP. The suggested added value of the combined treatments is emphasized in the improvement of chest expansion and spinal mobility.

Funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

CLINICAL RELEVANCE

- Both Kendall Exercises and Global Postural Reeducation could be clinically used to improve FHP with consideration of the available treatment time.
- The combined effect of Global postural reeducation with Kendall exercises do not have superior effect to Kendall exercise except for improving chest mobility, and spinal mobility in patients with FHP.
- Kendall exercises is proven to be a successful treatment for FHP.

REFERENCES

- Ackelman, B. H., & Lindgren, U. (2002). Validity and reliability of a modified version of the Neck Disability Index. *Journal of Rehabilitation Medicine*, 34(6), 284–287. <https://doi.org/10.1080/165019702760390383>
- Alowa, Z., & Elsayed, W. (2021). The impact of forward head posture on the electromyographic activity of the spinal muscles. *Journal of Taibah University Medical Sciences*, 16(2), 224–230. <https://doi.org/10.1016/J.JTUMED.2020.10.021>
- Audette, I., Dumas, J. P., Côté, J. N., & De Serres, S. J. (2010). Validity and between-day reliability of the cervical range of motion (CROM) device. *Journal of Orthopaedic and Sports Physical Therapy*, 40(5), 318–323. <https://doi.org/10.2519/JOSPT.2010.3180/ASSET/IMAGES/LARGE/JOSPT-318-FIG001.JPEG>

- Bennett, S., Siritaratiwat, W., Tanrangka, N., Bennett, M. J., & Kanpittaya, J. (2021). Effectiveness of the manual diaphragmatic stretching technique on respiratory function in cerebral palsy: A randomised controlled trial. *Respiratory Medicine*, *184*, 106443. <https://doi.org/10.1016/j.rmed.2021.106443>
- Blottner, D., Huang, Y., Trautmann, G., & Sun, L. (2019). The fascia: Continuum linking bone and myofascial bag for global and local body movement control on Earth and in Space. A scoping review. *REACH*, *14–15*, 100030. <https://doi.org/10.1016/J.REACH.2019.100030>
- Bonetti, F., Curti, S., Mattioli, S., Mugnai, R., Vanti, C., Violante, F. S., & Pillastrini, P. (2010). Effectiveness of a “Global Postural Reeducation” program for persistent Low Back Pain: a non-randomized controlled trial. <https://doi.org/10.1186/1471-2474-11-285>
- Bonney, R. A., & Corlett, E. N. (2002). Head posture and loading of the cervical spine. *Applied Ergonomics*, *33*(5), 415–417. [https://doi.org/10.1016/S0003-6870\(02\)00036-4](https://doi.org/10.1016/S0003-6870(02)00036-4)
- Bordoni, B., & Myers, T. (2020). A Review of the Theoretical Fascial Models: Biotensegrity, Fascintegrity, and Myofascial Chains. *Cureus*, *12*(2). <https://doi.org/10.7759/CUREUS.7092>
- Cagnie, B., Dickx, N., Peeters, I., Tuytens, J., Achten, E., Cambier, D., & Danneels, L. (2008). The use of functional MRI to evaluate cervical flexor activity during different cervical flexion exercises. *Journal of Applied Physiology*, *104*(1), 230–235. <https://doi.org/10.1152/JAPPLPHYSIOL.00918.2007>
- Carrasco-Lopez, V., & Medina-Porqueres, I. (2016). *Intervention using global postural reeducation method in patients with musculoskeletal diseases: systematic review*. *1*(2).
- Cavalcanti, I. F., Antonino, G. B., Do Monte-Silva, K. K., Guerino, M. R., Ferreira, A. P. D. L., & De Araújo, M. D. G. R. (2020). Global Postural Re-education in non-specific neck and low back pain treatment: A pilot study. *Journal of Back and Musculoskeletal Rehabilitation*, *33*(5), 823–828. <https://doi.org/10.3233/BMR-181371>
- Chang, M. C., Choo, Y. J., Hong, K., Boudier-Revéret, M., & Yang, S. (2023). Treatment of Upper Crossed Syndrome: A Narrative Systematic Review. *Healthcare (Basel, Switzerland)*, *11*(16). <https://doi.org/10.3390/HEALTHCARE11162328>
- Charmant, J., & Contributor. (2021). *Kinovea (0.9.5)*. Computer Software. <https://www.kinovea.org/>
- Cunha, A. C. V., Burke, T. N., França, F. J. R., & Marques, A. P. (2008). Effect of global posture reeducation and of static stretching on pain, range of motion, and quality of life in women with chronic neck pain: A randomized clinical trial. *Clinics*, *63*(6), 763–770. <https://doi.org/10.1590/S1807-59322008000600010>
- De Amorim, C. S. M., Gracitelli, M. E. C. M., Marques, A. A. A. P., Dos Santos Alves, V. L., Amorim, C., Gracitelli, M. E. C. M., Marques, A. A. A. P., & Alves, V. (2014). Effectiveness of global postural reeducation compared to segmental exercises on function, pain, and quality of life of patients with scapular dyskinesis associated with neck pain: A preliminary clinical trial. *Journal of Manipulative and Physiological Therapeutics*, *37*(6), 441–447. <https://doi.org/10.1016/j.jmpt.2013.08.011>
- De-La-Llave-Rincón, A. L., Domingo, P. C., Fernández-De-Las-Peñas, C., & Cleland, J. A. (2009). Increased Forward Head Posture and Restricted Cervical Range of Motion in Patients With Carpal Tunnel Syndrome. <https://doi.org/10.2519/Jospt.2009.3058>, *39*(9), 658–664. <https://doi.org/10.2519/JOSPT.2009.3058>

- Dimitrova, E., & Rohleva, M. (2014). Global postural reeducation in the treatment of postural impairments. *Resaarch in Kinesiology*, 4(1), 72–75.
- Fathollahnejad, K., Letafatkar, A., & Hadadnezhad, M. (2019). The effect of manual therapy and stabilizing exercises on forward head and rounded shoulder postures: A six-week intervention with a one-month follow-up study. *BMC Musculoskeletal Disorders*, 20(1), 1–8. <https://doi.org/10.1186/S12891-019-2438-Y/TABLES/5>
- Faul, F., Erdfelder, E., Lang, A. G., & Buchner, A. (2007). G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*, 39(2), 175–191. <https://doi.org/10.3758/BF03193146/METRICS>
- Fercho, J., Krakowiak, M., Yuser, R., Szmuda, T., Zieliński, P., Szarek, D., & Miękiśiak, G. (2023). Kinematic Analysis of the Forward Head Posture Associated with Smartphone Use. *Symmetry* 2023, Vol. 15, Page 667, 15(3), 667. <https://doi.org/10.3390/SYM15030667>
- Fernández-De-Las-Peñas, C., Alonso-Blanco, C., Morales-Cabezas, M., & Miangolarra-Page, J. C. (2005). Two exercise interventions for the management of patients with ankylosing spondylitis: A randomized controlled trial. *American Journal of Physical Medicine and Rehabilitation*, 84(6), 407–419. <https://doi.org/10.1097/01.phm.0000163862.89217.fe>
- Ferreira, G. E., Barreto, R. G. P., Robinson, C. C., Plentz, R. D. M., & Silva, M. F. (2016). Global Postural Reeducation for patients with musculoskeletal conditions: A systematic review of randomized controlled trials. In *Brazilian Journal of Physical Therapy* (Vol. 20, Issue 3, pp. 194–205). Revista Brasileira de Fisioterapia. <https://doi.org/10.1590/bjpt-rbf.2014.0153>
- Findley, T., Chaudhry, H., & Dhar, S. (2015). Transmission of muscle force to fascia during exercise. *Journal of Bodywork and Movement Therapies*, 19(1), 119–123. <https://doi.org/10.1016/J.JBMT.2014.08.010>
- França, F. R., Burke, T. N., Caffaro, R. R., Ramos, L. A., & Marques, A. P. (2012). Effects of muscular stretching and segmental stabilization on functional disability and pain in patients with chronic low back pain: A randomized, controlled trial. *Journal of Manipulative and Physiological Therapeutics*, 35(4), 279–285. <https://doi.org/10.1016/J.JMPT.2012.04.012>
- Fukaya, T., Konrad, A., Sato, S., Kiyono, R., Yahata, K., Yasaka, K., Onuma, R., Yoshida, R., & Nakamura, M. (2022). Comparison Between Contract–Relax Stretching and Antagonist Contract–Relax Stretching on Gastrocnemius Medialis Passive Properties. *Frontiers in Physiology*, 12, 764792. <https://doi.org/10.3389/FPHYS.2021.764792/BIBTEX>
- Guo, G., Zhang, Y., Lin, D., Chen, Z., Yan, Q., Gao, F., Wu, Y., He, J., Chen, D., Xie, Z., Huang, F., & Zhang, S. (2023). Correlation of finger-to-floor distance with the spinal mobility, spinal function indices and initial determination of its optimal cutoff value: a multicentre case–control study. *Frontiers in Medicine*, 10, 1135748. <https://doi.org/10.3389/FMED.2023.1135748>
- Harman, K., Hubley-Kozey, C. L., & Butler, H. (2005). Effectiveness of an exercise program to improve forward head posture in normal adults: A randomized, controlled 10-week trial. *Journal of Manual and Manipulative Therapy*, 13(3), 163–176. <https://doi.org/10.1179/106698105790824888>

- Hecimovich, M. D., & Hebert, J. J. (2016). Reliability and concurrent validity of an alternative method of lateral lumbar range of motion in athletes. *South African Journal of Sports Medicine*, 28(1), 23–26. <https://doi.org/10.17159/2078-516X/2016/V28I1A745>
- Heydari, Z., Sheikhhoseini, R., Shahrbanian, S., & Piri, H. (2022). Establishing minimal clinically important difference for effectiveness of corrective exercises on craniovertebral and shoulder angles among students with forward head posture: a clinical trial study. *BMC Pediatrics*, 22(1). <https://doi.org/10.1186/S12887-022-03300-7>
- Hindle, K., Whitcomb, T., Briggs, W., & Hong, J. (2012). Proprioceptive Neuromuscular Facilitation (PNF): Its Mechanisms and Effects on Range of Motion and Muscular Function. *Journal of Human Kinetics*, 31(1), 105. <https://doi.org/10.2478/V10078-012-0011-Y>
- Huijing, P. A. (2009). Epimuscular myofascial force transmission: A historical review and implications for new research. International society of biomechanics Muybridge award lecture, Taipei, 2007. *Journal of Biomechanics*, 42(1), 9–21. <https://doi.org/10.1016/J.JBIOMECH.2008.09.027>
- Im, B., Kim, Y., Chung, Y., & Hwang, S. (2016). Effects of scapular stabilization exercise on neck posture and muscle activation in individuals with neck pain and forward head posture. *Journal of Physical Therapy Science*, 28(3), 951. <https://doi.org/10.1589/JPTS.28.951>
- Jensen, M. P., Chen, C., & Brugger, A. M. (2003). Interpretation of visual analog scale ratings and change scores: A reanalysis of two clinical trials of postoperative pain. *Journal of Pain*, 4(7), 407–414. [https://doi.org/10.1016/S1526-5900\(03\)00716-8](https://doi.org/10.1016/S1526-5900(03)00716-8)
- Jørgensen, R., Ris, I., Juhl, C., Falla, D., & Juul-Kristensen, B. (2017). Responsiveness of clinical tests for people with neck pain. *BMC Musculoskeletal Disorders*, 18(1). <https://doi.org/10.1186/S12891-017-1918-1>
- Jung, S. I., Lee, N. K., Kang, K. W., Kim, K., & Lee, D. Y. (2016). The effect of smartphone usage time on posture and respiratory function. *Journal of Physical Therapy Science*, 28(1), 186. <https://doi.org/10.1589/JPTS.28.186>
- Karthik, V., Arulpragasame, S., Felix, A., & Parkavi, K. (2022). Prevalence Of Forward Head Posture And Its Association With Gender, BMI And Neck Pain Among College Going Students – A Cross Sectional Study. *Journal of Positive School Psychology*, 6(9), 5084–5090. <https://journalppw.com/index.php/jpsp/article/view/13487>
- Kendall, McCreary, Provance, Rodgers, & Romani. (2005). Muscles, Testing, and Function: With Posture and Pain. In *Muscles: testing and function with posture and pain* (5th ed.). Lippincott Williams & Wilkins.
- Khan, A., Khan, Z., Bhati, P., & Hussain, M. E. (2020). Influence of Forward Head Posture on Cervicocephalic Kinesthesia and Electromyographic Activity of Neck Musculature in Asymptomatic Individuals. *Journal of Chiropractic Medicine*, 19(4), 230–240. <https://doi.org/10.1016/J.JCM.2020.07.002>
- Khayatzaeh, S., Kalmanson, O. A., Schuit, D., Havey, R. M., Voronov, L. I., Ghanayem, A. J., & Patwardhan, A. G. (2017). Cervical Spine Muscle-Tendon Unit Length Differences Between Neutral and Forward Head Postures: Biomechanical Study Using Human Cadaveric Specimens. *Physical Therapy*, 97(7), 756–766. <https://doi.org/10.1093/PTJ/PZX040>

- Kim, D. H., Kim, C. J., & Son, S. M. (2018). Neck Pain in Adults with Forward Head Posture: Effects of Craniovertebral Angle and Cervical Range of Motion. *Osong Public Health and Research Perspectives*, 9(6), 309. <https://doi.org/10.24171/J.PHRP.2018.9.6.04>
- Kim, K.-H., Kim, S.-G., & Hwangbo, G. (2015). The effects of horse-riding simulator exercise and kendall exercise on the forward head posture. *Journal of Physical Therapy Science*, 27(4), 1125–1127. <https://doi.org/10.1589/jpts.27.1125>
- Kinjal Bagthariya, M. P. T., & Kakkad, A. (2024). “A Corelation Between Craniovertebral (CV) Angle, Cervical Range of Motion and Deep Neck Flexor Muscle Endurance in Young Adults with Neck Pain.” *Journal of Chemical Health Risks*, 14(2), 952–959. <https://doi.org/10.52783/JCHR.V14.I2.3553>
- Kong, Y. S., Kim, Y. M., & Shim, J. M. (2017). The effect of modified cervical exercise on smartphone users with forward head posture. *Journal of Physical Therapy Science*, 29(2), 328. <https://doi.org/10.1589/JPTS.29.328>
- Kose, E., Baylan, H., Karahan, H., Cetin Ekerbicer, H., & Baylan, H. (2022). The Distribution and The Related Factors of Forward Head Posture Among Medical Students. *Konuralp Medical Journal*, 14(2), 304–308. <https://doi.org/10.18521/ktd.952182>
- Koseki, T., Kakizaki, F., Hayashi, S., Nishida, N., & Itoh, M. (2019). Effect of forward head posture on thoracic shape and respiratory function. *Journal of Physical Therapy Science*, 31(1), 63. <https://doi.org/10.1589/JPTS.31.63>
- Lee, N. K., Jung, S. I., Lee, D. Y., & Kang, K. W. (2017). Effects of Exercise on Cervical Angle and Respiratory Function in Smartphone Users. *Osong Public Health and Research Perspectives*, 8(4), 271. <https://doi.org/10.24171/J.PHRP.2017.8.4.07>
- Lee, S. M., Lee, C. H., O’Sullivan, D., Jung, J. H., & Park, J. J. (2016). Clinical effectiveness of a Pilates treatment for forward head posture. *Journal of Physical Therapy Science*, 28(7), 2009. <https://doi.org/10.1589/JPTS.28.2009>
- Lin, G., Wang, W., & Wilkinson, T. (2022). Changes in deep neck muscle length from the neutral to forward head posture. A cadaveric study using Thiel cadavers. *Clinical Anatomy (New York, N.Y.)*, 35(3), 332–339. <https://doi.org/10.1002/CA.23834>
- Macdelilid, J. C., Walton, D. M., Avery, S., Blanchard, A., Etruw, E., Mcalpine, C., & Goldsmith, C. H. (2009). Measurement properties of the neck disability index: a systematic review. *The Journal of Orthopaedic and Sports Physical Therapy*, 39(5), 400–416. <https://doi.org/10.2519/JOSPT.2009.2930>
- Matsutani, L. A., Sousa do Espírito Santo, A. de, Ciscato, M., Yuan, S. L. K., & Marques, A. P. (2023). Global posture reeducation compared with segmental muscle stretching exercises in the treatment of fibromyalgia: a randomized controlled trial. *Trials*, 24(1), 1–13. <https://doi.org/10.1186/S13063-023-07422-W/TABLES/6>
- Mendes Fernandes, T., Méndez-Sánchez, R., Puente-González, A. S., Martín-Vallejo, F. J., Falla, D., & Vila-Chã, C. (2023). A randomized controlled trial on the effects of “Global Postural Re-education” versus neck specific exercise on pain, disability, postural control, and neuromuscular features in women with chronic non-specific neck pain. *European Journal of Physical and Rehabilitation Medicine*, 59(1), 42. <https://doi.org/10.23736/S1973-9087.22.07554-2>
- Mendes-Fernandes, T., Puente-González, A. S., Márquez-Vera, M. A., Vila-Chã, C., & Méndez-Sánchez, R. (2021). Effects of Global Postural Reeducation versus Specific Therapeutic Neck Exercises on Pain, Disability, Postural Control, and Neuromuscular

- Efficiency in Women with Chronic Nonspecific Neck Pain: Study Protocol for a Randomized, Parallel, Clinical Trial. *International Journal of Environmental Research and Public Health*, 18(20), 10704. <https://doi.org/10.3390/IJERPH182010704>
- Mohan, V., Dzulkifli, N. H., Justine, M., Haron, R., Leonard Joseph, H., & Rathinam, C. (2012). Intrarater reliability of chest expansion using cloth tape measure technique. *Bangladesh Journal of Medical Science*, 11(4), 307–311. <https://doi.org/10.3329/BJMS.V11i4.12602>
- Moreno, M. A., Catai, A. M., Teodori, R. M., Borges, B. L. A., De Cesar, M. C., & Da Silva, E. (2007). Effect of a muscle stretching program using the Global Postural Reeducation method on respiratory muscle strength and thoracoabdominal mobility of sedentary young males. *Jornal Brasileiro de Pneumologia*, 33(6), 679–686. <https://doi.org/10.1590/S1806-37132007000600011>
- Moreno, M. A., Catai, A. M., Teodori, R. M., Borges, B. LA, De Castro Cesar, M., & Da Silva, E. (2007). Effect of a muscle stretching program using the Global Postural Re-education method on respiratory muscle strength and thoracoabdominal mobility of sedentary young males*. *J Bras Pneumol*, 33(6), 679–686.
- Moustafa, I. M., Youssef, A., Ahbouch, A., Tamim, M., & Harrison, D. E. (2020). Is forward head posture relevant to autonomic nervous system function and cervical sensorimotor control? Cross sectional study. *Gait & Posture*, 77, 29–35. <https://doi.org/10.1016/j.gaitpost.2020.01.004>
- Mun Cheung Lau, H., Tai Wing Chiu, T., St, Mp., & Lam, T.-H. (2010). Measurement of craniovertebral angle with Electronic Head Posture Instrument: Criterion validity. *Journal of Rehabilitation Research & Development*, 47(9), 911–918. <https://doi.org/10.1682/JRRD.2010.01.0001>
- Najwatul, N., Beng Gan, K., Jian Sia, Z., Abdul Karim, Z., & Kaur Ajit Singh, D. (2016). CONCURRENT VALIDITY OF CHEST WALL EXPANSION BETWEEN CLOTH TAPE MEASURE, ELECTROMAGNETIC AND LASER DISPLACEMENT SENSORS. *Malaysian Journal of Industrial Technology*, 1(2). www.mitec.unikl.edu.my/mjit
- Oliveri, M., Caltagirone, C., Loriga, R., Pompa, M. N., Versace, V., & Souchard, P. (2012). Fast increase of motor cortical inhibition following postural changes in healthy subjects. *Neuroscience Letters*, 530(1), 7–11. <https://doi.org/10.1016/j.neulet.2012.09.031>
- Page, S. J., & Persch, A. C. (2013). Recruitment, Retention, and Blinding in Clinical Trials. *The American Journal of Occupational Therapy*, 67(2), 154. <https://doi.org/10.5014/AJOT.2013.006197>
- Perret, C., Poiraudreau, S., Fermanian, J., Lefèvre Colau, M. M., Mayoux Benhamou, M. A., & Revel, M. (2001). Validity, reliability, and responsiveness of the fingertip-to-floor test. *Archives of Physical Medicine and Rehabilitation*, 82(11), 1566–1570. <https://doi.org/10.1053/APMR.2001.26064>
- Pillastrini, P., Banchelli, F., Guccione, A., Di Ciaccio, E., Violante, F. S., Brugnettoni, M., & Vanti, C. (2018). Global Postural Reeducation in patients with chronic nonspecific neck pain: cross-over analysis of a randomized controlled trial. *La Medicina Del Lavoro*, 109(1), 16. <https://doi.org/10.23749/MDL.V109I1.6677>
- Pt, I. A. Y., Dunning, J., Butts Pt, R., Pt, M., & Pt, J. A. C. (2018). Reliability, construct validity, and responsiveness of the neck disability index and numeric pain rating scale in patients with mechanical neck pain without upper extremity symptoms. *Physiotherapy*

- Theory and Practice*, 35(12), 1328–1335.
<https://doi.org/10.1080/09593985.2018.1471763>
- Puig-Diví, A., Escalona-Marfil, C., Padullés-Riu, J. M., Busquets, A., Padullés-Chando, X., & Marcos-Ruiz, D. (2019). Validity and reliability of the Kinovea program in obtaining angles and distances using coordinates in 4 perspectives. *PLOS ONE*, 14(6), e0216448.
<https://doi.org/10.1371/JOURNAL.PONE.0216448>
- Rahul S, Anitha A, Kamalakannan M, & Ramana K. (2024). Effectiveness of Kendall Exercise for Forward Head Posture among IT Workers. *Indian Journal of Physiotherapy & Occupational Therapy - An International Journal*, 18, 743–748.
<https://doi.org/10.37506/dcmj2p97>
- Reddy, R. S., Alahmari, K. A., Silvian, P. S., Ahmad, I. A., Kakarparthi, V. N., & Rengaramanujam, K. (2019). Reliability of Chest Wall Mobility and Its Correlation with Lung Functions in Healthy Nonsmokers, Healthy Smokers, and Patients with COPD. *Canadian Respiratory Journal*, 2019. <https://doi.org/10.1155/2019/5175949>
- Rocha, L. S. de O., Mineshita, L. N. H., Sobral, L. L., Magno, L. D., Santos, M. C. de S., Rocha, R. S. B., Rocha, L. S. de O., Mineshita, L. N. H., Sobral, L. L., Magno, L. D., Santos, M. C. de S., & Rocha, R. S. B. (2018). Influence of global postural reeducation method on respiratory muscle strength and parkinsonian quality of life. *Manual Therapy, Posturology & Rehabilitation Journal*, 15, 0–0.
<https://doi.org/10.17784/MTPREHABJOURNAL.2017.15.504>
- Rosário, J. L. P. do, Nakashima, I. Y., Rizopoulos, K., Kostopoulos, D., & Marques, A. P. (2012). Improving posture: Comparing Segmental Stretch and Muscular Chains Therapy. *Clinical Chiropractic*, 15(3–4), 121–128.
<https://doi.org/10.1016/j.clch.2012.10.039>
- Ruivo, R. M., Pezarat-Correia, P., & Carita, A. I. (2017). Effects of a Resistance and Stretching Training Program on Forward Head and Protracted Shoulder Posture in Adolescents. *Journal of Manipulative and Physiological Therapeutics*, 40(1), 1–10.
<https://doi.org/10.1016/J.JMPT.2016.10.005>
- Sarraf, F., & Varmazyar, S. (2022). Comparing the effect of the posture of using smartphones on head and neck angles among college students. *Ergonomics*, 65(12), 1631–1638. <https://doi.org/10.1080/00140139.2022.2047229>
- Schwertner, D. S., Oliveira, R. A. N. da S., Beltrame, T. S., Capistrano, R., & Alexandre, J. M. (2018). Questionnaire on body awareness of postural habits in young people: construction and validation. *Fisioterapia Em Movimento*, 31(0), e003116.
<https://doi.org/10.1590/1980-5918.031.AO16>
- Sheikhoseini, R., Shahrbanian, S., Sayyadi, P., & O’Sullivan, K. (2018). Effectiveness of Therapeutic Exercise on Forward Head Posture: A Systematic Review and Meta-analysis. *Journal of Manipulative and Physiological Therapeutics*, 41(6), 530–539.
<https://doi.org/10.1016/J.JMPT.2018.02.002>
- Siemonsma, P. C., Stuive, I., Roorda, L. D., Vollebregt, J. A., Walker, M. F., Lankhorst, G. J., & Lettinga, A. T. (2013). *Cognitive Treatment of Illness Perceptions in Patients With Chronic Low Back Pain: A Randomized Controlled Trial*.
<https://academic.oup.com/ptj/article/93/4/435/2735290>
- Silva, E. M., Andrade, S. C., Vilar, M. J., Silva, M., Andrade, S. C., & Vilar, M. J. (2012). Evaluation of the effects of Global Postural Reeducation in patients with ankylosing

- spondylitis. *Rheumatology International*, 32(7), 2155–2163.
<https://doi.org/10.1007/s00296-011-1938-3>
- Singla, D., Veqar, Z., & Hussain, M. E. (2017). Photogrammetric Assessment of Upper Body Posture Using Postural Angles: A Literature Review. *Journal of Chiropractic Medicine*, 16(2), 131–138. <https://doi.org/10.1016/j.jcm.2017.01.005>
- Souchard, P. E., Meli, O., Sgamma, D., Pastor, I., Korell, M., & Michel, B. (2011). *Rééducation posturale globale RPG - La méthode*. Elsevier Masson.
- Stecco, A., Giordani, F., Fede, C., Pirri, C., De Caro, R., & Stecco, C. (2023). From Muscle to the Myofascial Unit: Current Evidence and Future Perspectives. *International Journal of Molecular Sciences*, 24(5), 24. <https://doi.org/10.3390/IJMS24054527>
- Tousignant, M., Smeesters, C., Breton, A. M., Breton, É., & Corriveau, H. (2006). Criterion validity study of the cervical range of motion (CROM) device for rotational range of motion on healthy adults. *Journal of Orthopaedic and Sports Physical Therapy*, 36(4), 242–248. <https://doi.org/10.2519/jospt.2006.36.4.242>
- Van Det, M. J., Meijerink, W. J. H. J., Hoff, C., Van Veelen, M. A., & Pierie, J. P. E. N. (2008). Ergonomic assessment of neck posture in the minimally invasive surgery suite during laparoscopic cholecystectomy. *Surgical Endoscopy and Other Interventional Techniques*, 22(11), 2421–2427. <https://doi.org/10.1007/S00464-008-0042-6>
- Vernon, H., & Mior, S. (1991). The Neck Disability Index: a study of reliability and validity. *Journal of Manipulative and Physiological Therapeutics*, 14(7), 409–415.
<http://europepmc.org/article/med/1834753>
- Wang, S.-F., Chai, H.-M., & Lu, T.-W. (2002). Comparison of Ranges of Cervical Motion Measured by Gravity-Based Goniometry and Ultrasound-Based Motion Analysis System. *FJPT*, 27(3), 124–130.
- Wilke, J., & Krause, F. (2019). Myofascial chains of the upper limb: A systematic review of anatomical studies. *Clinical Anatomy*, 32(7), 934–940.
<https://doi.org/10.1002/CA.23424>
- Wilke, J., Krause, F., Vogt, L., & Banzer, W. (2016). What is evidence-based about myofascial chains: A systematic review. *Archives of Physical Medicine and Rehabilitation*, 97(3), 454–461. <https://doi.org/10.1016/J.APMR.2015.07.023>
- Wilke, J., Schleip, R., Yucesoy, C. A., & Banzer, W. (2018). Not merely a protective packing organ? A review of fascia and its force transmission capacity. *Journal of Applied Physiology*, 124(1), 234–244. <https://doi.org/10.1152/JAPPLPHYSIOL.00565.2017>
- Wilke, J., Vogt, L., Niederer, D., & Banzer, W. (2017). Is remote stretching based on myofascial chains as effective as local exercise? A randomised-controlled trial. *Journal of Sports Sciences*, 35(20), 2021–2027.
<https://doi.org/10.1080/02640414.2016.1251606>
- Williams, M. A., McCarthy, C. J., Chorti, A., Cooke, M. W., & Gates, S. (2010). A Systematic Review of Reliability and Validity Studies of Methods for Measuring Active and Passive Cervical Range of Motion. *Journal of Manipulative and Physiological Therapeutics*, 33(2), 138–155. <https://doi.org/10.1016/J.JMPT.2009.12.009>
- Xu, L., Hwang, B., & Kim, T. (2019). The Effect of Postural Correction and Visual Feedback on Muscle Activity and Head Position Change During Overhead Arm Lift Test in Subjects with Forward Head Posture. *The Journal of Korean Physical Therapy*, 31(3), 151–156.
<https://doi.org/10.18857/JKPT.2019.31.3.151>

- Yagci, G., Ayhan, C., & Yakut, Y. (2018). Effectiveness of basic body awareness therapy in adolescents with idiopathic scoliosis: A randomized controlled study¹. *Journal of Back and Musculoskeletal Rehabilitation*, 31(4), 693–701. <https://doi.org/10.3233/BMR-170868>
- Yip, C. H. T., Chiu, T. T. W., & Poon, A. T. K. (2008). The relationship between head posture and severity and disability of patients with neck pain. *Manual Therapy*, 13(2), 148–154. <https://doi.org/10.1016/J.MATH.2006.11.002>

TABLES

Table 1: Exercises, dose, and progression of Kendall exercises.

Exercise	Dose	Description and Progression
Strengthening of Deep Cervical Flexors	5 sets of 10 repetitions, each set interrupted by 30 seconds break.	Chin tucks in supine lying with the head in contact with the floor. Progression from the third week by lifting the head off the floor in a tucked position and holding it for 2-8 seconds.
Stretching the cervical extensors	Five repetitions were held for 60 seconds each and followed by 60 seconds break.	a chin drop-in sitting progresses by chin drop with hand assistance by placing both hands on the occipital area.
Strengthening shoulder retraction	Five sets of 10 repetitions, each set interrupted by 30 seconds break.	Shoulder retraction from standing by holding a TheraBand moving scapular blades inwards. Progression to prone lying weight-free then using the weight of 1 kg.
Stretching the pectoralis muscle	Five repetitions were held for 60 seconds each and followed by 60 seconds break.	Unilateral Pectoral Stretch using the door frame or wall. Progresses to bilateral stretch and increases the force of stretch as tolerated.

Table 2: GPR postures, sequence, and description as used in the current study

Shoulder and Hip Angles	Postures	Engaged Chains	Application time and duration	Description
Open Coxofemoral (Hip Joint) and Shoulders adducted	Frog on the Ground	anterior chain, respiratory chain, anterior arm, upper scapular chains, and anteromedial and lateral hip chains.	Week 1 and 2 applied for 15 minutes.	Supine position, neck, and shoulders retracted, arms adducted from 45° to 0° throughout the sessions, elbow extended, abdomen retracted, pelvis posteriorly tilted, hips abducted, and externally rotated, knees flexed, and feet faced.
	Standing against the wall	Anterior chain, respiratory chain, anterior arm, upper scapular chains, and anteromedial and lateral hip chains. This also improves balance and proprioception.	Week 3 and 4 applied for 15 minutes.	Standing against the wall, focus on the alignment of the spine, chin retracted, shoulders retracted, arms adducted from 45° to 0° over the course of the sessions, elbow extended, abdomen retracted, pelvis posteriorly tilted, hips extended, and knees bent to extension as tolerated.
Closed Coxofemoral (Hip Joint) and Shoulders adducted	Frog in the Air	Posterior chain, respiratory chain; anterointernal of the shoulder, anterior arm, and side chain of the hip.	Week 1 and 2 applied for 15 minutes.	Supine, chin retracted, shoulders posteriorly tilted, elbows extended, abdomen retracted, pelvis posteriorly tilted, hips flexed, knees flexed and the ankles resting on the wall with the arms started at 45° to 0°.
	Sitting	Posterior chain, respiratory chain; anterointernal of the shoulder; scapular chain, anterior arm, and side chain of the hip.	Week 4 applied for 15 minutes.	Sitting, focus on the alignment of the spine, chin in, shoulders retracted as tolerated, abdomen retracted, knees bent, arms adducted 45° to 0°.

Table 3. Comparison of subject characteristics between groups A and B:

	Group A	Group B	MD	t- value	p-value
Age (years)	21.09 ± 3.10	22.71 ± 3.72	-1.62	-1.55	0.13
Weight (kg)	77.91 ± 14.72	75.88 ± 10.47	2.03	0.52	0.61
Height (cm)	167.41 ± 8.32	170.09 ± 11.17	-2.68	-0.89	0.37
BMI (kg/m²)	27.88 ± 5.09	26.41 ± 4.26	1.47	1.03	0.31
Sex, n (%)					
Female	12 (54.5%)	11 (52%)		(χ ² = 0.02)	0.88
Male	10 (45.5%)	10 (48%)			

SD, Standard deviation; t, unpaired t value; χ², Chi-squared value; p-value, Level of significance

Table 4 Mean CVA, GA, and SA, CROM, Chest expansion (axillary and xyphoid), and fingers-to-floor test pre and post-treatment of groups A and B.

Outcome Measures	Group A		Group B		MD (95% CI)	p-value	Effect size
	Mean \pm SD	Mean \pm SD	Mean \pm SD	Mean \pm SD			
CVA (degrees)	Pretreatment	44.94 \pm 2.13	44.87 \pm 2.59		0.07 (-1.39: 1.52)	0.92	0.17
	Post-treatment	51.83 \pm 3.32	51.13 \pm 4.61		0.7 (-1.76: 3.17)	0.56	
	MD (95% CI)	-6.89 (-8.60: -5.18)	-6.26 (-8: -4.5)				
	% of change	15.33 <i>p</i> = 0.001*	13.95 <i>p</i> = 0.001*				
GA (degrees)	Pretreatment	17.35 \pm 6.97	17.99 \pm 3.51		0.64 (-4.06-2.79)	0.711	0.006
	Post-treatment	17.79 \pm 5.45	18.59 \pm 5.04		0.8 (-3.93-2.33)	0.609	
	MD (95% CI)	0.62 (-2.18-1.31)	0.6 (-2.39-1.19)				
	% of change	3.57 <i>P</i> = 0.617	3.34 <i>P</i> = 0.500				
SA (degrees)	Pretreatment	49.57 \pm 13.11	54.33 \pm 15.48		-4.76 (-13.58: 4.06)	0.28	0.22
	Post-treatment	53.25 \pm 11.34	50.49 \pm 13.93		2.76 (-5.04: 10.57)	0.48	
	MD (95% CI)	-3.68 (-9.55: 2.19)	3.84 (-2.16: 9.85)				
	% of change	7.42 <i>p</i> = 0.21	7.07 <i>p</i> = 0.20				
Flexion (degrees)	Pretreatment	51.39 \pm 10.56	52.22 \pm 11.82		-0.83 (-7.73: 6.07)	0.81	0.05
	Post-treatment	57.80 \pm 10.09	58.22 \pm 7.42		-0.42 (-5.90: 5.06)	0.87	
	MD (95% CI)	-6.41 (-10.60: -2.2)	-6 (-10.29: -1.71)				
	% of change	12.47 <i>p</i> = 0.004*	11.49 <i>p</i> = 0.007*				
Extension (degrees)	Pretreatment	52.32 \pm 7.79	55.79 \pm 9.41		-3.47 (-8.78: 1.83)	0.19	0.16
	Post-treatment	60.30 \pm 11.37	62.11 \pm 10.46		-1.81 (-8.55: 4.93)	0.59	
	MD (95% CI)	-7.98 (-12.77: -3.20)	-6.32 (-11.22: -1.42)				
	% of change	15.25 <i>p</i> = 0.002*	11.33 <i>p</i> = 0.01*				
Right side bending (degrees)	Pretreatment	40.50 \pm 6.37	39.92 \pm 6.96		0.58 (-3.53: 4.69)	0.77	0.03
	Post-treatment	46.06 \pm 8.55	45.87 \pm 6.15		0.19 (-4.42: 4.80)	0.93	
	MD (95% CI)	-5.56 (-8.86: -2.26)	-5.95 (-9.33: -2.57)				
	% of change	13.73 <i>p</i> = 0.002*	14.90 <i>p</i> = 0.001*				
	Pretreatment	38.69 \pm 6.99	40.67 \pm 8.59		-1.98 (-6.78: 2.84)	0.41	

Left side bending (degrees)	Post-treatment	44.09 ± 8.32	46 ± 8.93	-1.91 (-7.22: 3.41)	0.47	0.22
	MD (95% CI)	-5.4 (-8.55: -2.24)	-5.33 (-8.56: -2.11)			
	% of change	13.96	13.11			
		<i>p</i> = 0.001*	<i>p</i> = 0.002*			
Right Rotation (degrees)	Pretreatment	58.45 ± 12.35	58.63 ± 10.43	-0.18 (-7.24: 6.88)	0.96	0.23
	Post-treatment	67.12 ± 6.59	65.35 ± 8.84	1.77 (-3.01: 5.56)	0.46	
	MD (95% CI)	-8.67 (-13.37: -3.96)	-6.72 (-11.53: -1.90)			
	% of change	14.83	11.46			
		<i>p</i> = 0.001*	<i>p</i> = 0.007*			
Left rotation (degrees)	Pretreatment	59.45 ± 8.11	57.08 ± 11.53	2.37 (-3.74: 8.49)	0.43	0.23
	Post-treatment	67.80 ± 9.33	65.83 ± 7.88	1.97 (-3.36: 7.31)	0.45	
	MD (95% CI)	-8.35 (-12.74: -3.96)	-8.75 (-13.24: -4.25)			
	% of change	14.05	15.33			
		<i>p</i> = 0.001*	<i>p</i> = 0.001*			
Axillary circumference (centimeter)	Pretreatment	99.36 ± 8.54	98.76 ± 7.89	0.6 (-4.47: 5.67)	0.81	0.76
	Post-treatment	105.05 ± 8.34	98.62 ± 8.56	6.43 (1.22: 11.63)	0.01	
	MD (95% CI)	-5.69 (-6.96: -4.41)	0.14 (-1.16: 1.44)			
	% of change	-5.73	0.14			
		<i>p</i> = 0.001*	<i>p</i> = 0.83			
Sternal circumference (centimeter)	Pretreatment	90.05 ± 8.78	91.47 ± 7.17	-1.42 (-6.38: 3.52)	0.56	0.80
	Post-treatment	97.09 ± 8.87	90.38 ± 7.77	6.71 (1.56: 11.86)	0.01	
	MD (95% CI)	-7.04 (-8.46: -5.63)	1.09 (-0.35: 2.54)			
	% of change	7.82	1.19			
		<i>p</i> = 0.001*	<i>p</i> = 0.13			
Finger to floor test (centimeter)	Pretreatment	9.97 ± 9.58	9.88 ± 9.99	0.09 (-5.93: 6.13)	0.97	0.75
	Post-treatment	2.29 ± 8.49	9.14 ± 9.61	-6.85 (-12.43: -1.27)	0.01	
	MD (95% CI)	7.68 (6.24: 9.12)	0.74 (-0.73: 2.21)			
	% of change	77.03	7.49			
		<i>p</i> = 0.001*	<i>p</i> = 0.32			

SD, Standard deviation; MD, Mean difference; CI, Confidence interval; p-value, Probability value.* indicate significance

Table 5 Median NDI pre- and post-treatment of groups A and B:

Outcome		Group A	Group B		
Measures		Median (IQR)	Median (IQR)	U-value	p-value
NDI (%)	Pretreatment	4 (4-2)	4 (4-4)	201	0.40
	Post-treatment	2 (4-2)	2 (4-2)	215.5	0.66
	Z-value	2.31	2.48		
		<i>p = 0.02*</i>	<i>p = 0.01*</i>		

IQR, Interquartile range; Z- value, Wilcoxon signed ranks test value; U- value, Mann-Whitney test value; p-value, Probability value. * indicate significance.

Journal Pre-proof

TABLES

Table 1: Exercises, dose, and progression of Kendall exercises.

Exercise	Dose	Description and Progression
Strengthening of Deep Cervical Flexors	5 sets of 10 repetitions, each set interrupted by 30 seconds break.	Chin tucks in supine lying with the head in contact with the floor. Progression from the third week by lifting the head off the floor in a tucked position and holding it for 2-8 seconds.
Stretching the cervical extensors	Five repetitions were held for 60 seconds each and followed by 60 seconds break.	a chin drop-in sitting progresses by chin drop with hand assistance by placing both hands on the occipital area.
Strengthening shoulder retraction	Five sets of 10 repetitions, each set interrupted by 30 seconds break.	Shoulder retraction from standing by holding a TheraBand moving scapular blades inwards. Progression to prone lying weight-free then using the weight of 1 kg.
Stretching the pectoralis muscle	Five repetitions were held for 60 seconds each and followed by 60 seconds break.	Unilateral Pectoral Stretch using the door frame or wall. Progresses to bilateral stretch and increases the force of stretch as tolerated.

Table 2: GPR postures, sequence, and description as used in the current study

Shoulder and Hip Angles	Postures	Engaged Chains	Application time and duration	Description
Open Coxofemoral (Hip Joint) and Shoulders adducted	Frog on the Ground	anterior chain, respiratory chain, anterior arm, upper scapular chains, and anteromedial and lateral hip chains.	Week 1 and 2 applied for 15 minutes.	Supine position, neck, and shoulders retracted, arms adducted from 45° to 0° throughout the sessions, elbow extended, abdomen retracted, pelvis posteriorly tilted, hips abducted, and externally rotated, knees flexed, and feet faced.
	Standing against the wall	Anterior chain, respiratory chain, anterior arm, upper scapular chains, and anteromedial and lateral hip chains. This also improves balance and proprioception.	Week 3 and 4 applied for 15 minutes.	Standing against the wall, focus on the alignment of the spine, chin retracted, shoulders retracted, arms adducted from 45° to 0° over the course of the sessions, elbow extended, abdomen retracted, pelvis posteriorly tilted, hips extended, and knees bent to extension as tolerated.
Closed Coxofemoral (Hip Joint) and Shoulders adducted	Frog in the Air	Posterior chain, respiratory chain; anterointernal of the shoulder, anterior arm, and side chain of the hip.	Week 1 and 2 applied for 15 minutes.	Supine, chin retracted, shoulders posteriorly tilted, elbows extended, abdomen retracted, pelvis posteriorly tilted, hips flexed, knees flexed and the ankles resting on the wall with the arms started at 45° to 0°.
	Sitting	Posterior chain, respiratory chain; anterointernal of the shoulder; scapular chain, anterior arm, and side chain of the hip.	Week 4 applied for 15 minutes.	Sitting, focus on the alignment of the spine, chin in, shoulders retracted as tolerated, abdomen retracted, knees bent, arms adducted 45° to 0°.

Table 3. Comparison of subject characteristics between groups A and B:

	Group A	Group B	MD	t- value	p-value
Age (years)	21.09 ± 3.10	22.71 ± 3.72	-1.62	-1.55	0.13
Weight (kg)	77.91 ± 14.72	75.88 ± 10.47	2.03	0.52	0.61
Height (cm)	167.41 ± 8.32	170.09 ± 11.17	-2.68	-0.89	0.37
BMI (kg/m²)	27.88 ± 5.09	26.41 ± 4.26	1.47	1.03	0.31
Sex, n (%)					
Female	12 (54.5%)	11 (52%)		(χ ² = 0.02)	0.88
Male	10 (45.5%)	10 (48%)			

SD, Standard deviation; t, unpaired t value; χ², Chi-squared value; p-value, Level of significance

Table 4 Mean CVA, GA, and SA, CROM, Chest expansion (axillary and xyphoid), and fingers-to-floor test pre and post-treatment of groups A and B.

Outcome Measures	Group A		Group B		MD (95% CI)	p-value	Effect size
	Mean \pm SD	Mean \pm SD	Mean \pm SD	Mean \pm SD			
CVA (degrees)	Pretreatment	44.94 \pm 2.13	44.87 \pm 2.59		0.07 (-1.39: 1.52)	0.92	0.17
	Post-treatment	51.83 \pm 3.32	51.13 \pm 4.61		0.7 (-1.76: 3.17)	0.56	
	MD (95% CI)	-6.89 (-8.60: -5.18)	-6.26 (-8: -4.5)				
	% of change	15.33 <i>p</i> = 0.001*	13.95 <i>p</i> = 0.001*				
GA (degrees)	Pretreatment	17.35 \pm 6.97	17.99 \pm 3.51		0.64 (-4.06-2.79)	0.711	0.006
	Post-treatment	17.79 \pm 5.45	18.59 \pm 5.04		0.8 (-3.93-2.33)	0.609	
	MD (95% CI)	0.62 (-2.18-1.31)	0.6 (-2.39-1.19)				
	% of change	3.57 <i>P</i> = 0.617	3.34 <i>P</i> = 0.500				
SA (degrees)	Pretreatment	49.57 \pm 13.11	54.33 \pm 15.48		-4.76 (-13.58: 4.06)	0.28	0.22
	Post-treatment	53.25 \pm 11.34	50.49 \pm 13.93		2.76 (-5.04: 10.57)	0.48	
	MD (95% CI)	-3.68 (-9.55: 2.19)	3.84 (-2.16: 9.85)				
	% of change	7.42 <i>p</i> = 0.21	7.07 <i>p</i> = 0.20				
Flexion (degrees)	Pretreatment	51.39 \pm 10.56	52.22 \pm 11.82		-0.83 (-7.73: 6.07)	0.81	0.05
	Post-treatment	57.80 \pm 10.09	58.22 \pm 7.42		-0.42 (-5.90: 5.06)	0.87	
	MD (95% CI)	-6.41 (-10.60: -2.2)	-6 (-10.29: -1.71)				
	% of change	12.47 <i>p</i> = 0.004*	11.49 <i>p</i> = 0.007*				
Extension (degrees)	Pretreatment	52.32 \pm 7.79	55.79 \pm 9.41		-3.47 (-8.78: 1.83)	0.19	0.16
	Post-treatment	60.30 \pm 11.37	62.11 \pm 10.46		-1.81 (-8.55: 4.93)	0.59	
	MD (95% CI)	-7.98 (-12.77: -3.20)	-6.32 (-11.22: -1.42)				
	% of change	15.25 <i>p</i> = 0.002*	11.33 <i>p</i> = 0.01*				
Right side bending (degrees)	Pretreatment	40.50 \pm 6.37	39.92 \pm 6.96		0.58 (-3.53: 4.69)	0.77	0.03
	Post-treatment	46.06 \pm 8.55	45.87 \pm 6.15		0.19 (-4.42: 4.80)	0.93	
	MD (95% CI)	-5.56 (-8.86: -2.26)	-5.95 (-9.33: -2.57)				
	% of change	13.73 <i>p</i> = 0.002*	14.90 <i>p</i> = 0.001*				
	Pretreatment	38.69 \pm 6.99	40.67 \pm 8.59		-1.98 (-6.78: 2.84)	0.41	

Left side bending (degrees)	Post-treatment	44.09 ± 8.32	46 ± 8.93	-1.91 (-7.22: 3.41)	0.47	0.22
	MD (95% CI)	-5.4 (-8.55: -2.24)	-5.33 (-8.56: -2.11)			
	% of change	13.96	13.11			
		<i>p</i> = 0.001*	<i>p</i> = 0.002*			
Right Rotation (degrees)	Pretreatment	58.45 ± 12.35	58.63 ± 10.43	-0.18 (-7.24: 6.88)	0.96	0.23
	Post-treatment	67.12 ± 6.59	65.35 ± 8.84	1.77 (-3.01: 5.56)	0.46	
	MD (95% CI)	-8.67 (-13.37: -3.96)	-6.72 (-11.53: -1.90)			
	% of change	14.83	11.46			
		<i>p</i> = 0.001*	<i>p</i> = 0.007*			
Left rotation (degrees)	Pretreatment	59.45 ± 8.11	57.08 ± 11.53	2.37 (-3.74: 8.49)	0.43	0.23
	Post-treatment	67.80 ± 9.33	65.83 ± 7.88	1.97 (-3.36: 7.31)	0.45	
	MD (95% CI)	-8.35 (-12.74: -3.96)	-8.75 (-13.24: -4.25)			
	% of change	14.05	15.33			
		<i>p</i> = 0.001*	<i>p</i> = 0.001*			
Axillary circumference (centimeter)	Pretreatment	99.36 ± 8.54	98.76 ± 7.89	0.6 (-4.47: 5.67)	0.81	0.76
	Post-treatment	105.05 ± 8.34	98.62 ± 8.56	6.43 (1.22: 11.63)	0.01	
	MD (95% CI)	-5.69 (-6.96: -4.41)	0.14 (-1.16: 1.44)			
	% of change	-5.73	0.14			
		<i>p</i> = 0.001*	<i>p</i> = 0.83			
Sternal circumference (centimeter)	Pretreatment	90.05 ± 8.78	91.47 ± 7.17	-1.42 (-6.38: 3.52)	0.56	0.80
	Post-treatment	97.09 ± 8.87	90.38 ± 7.77	6.71 (1.56: 11.86)	0.01	
	MD (95% CI)	-7.04 (-8.46: -5.63)	1.09 (-0.35: 2.54)			
	% of change	7.82	1.19			
		<i>p</i> = 0.001*	<i>p</i> = 0.13			
Finger to floor test (centimeter)	Pretreatment	9.97 ± 9.58	9.88 ± 9.99	0.09 (-5.93: 6.13)	0.97	0.75
	Post-treatment	2.29 ± 8.49	9.14 ± 9.61	-6.85 (-12.43: -1.27)	0.01	
	MD (95% CI)	7.68 (6.24: 9.12)	0.74 (-0.73: 2.21)			
	% of change	77.03	7.49			
		<i>p</i> = 0.001*	<i>p</i> = 0.32			

SD, Standard deviation; MD, Mean difference; CI, Confidence interval; p-value, Probability value.* indicate significance

Table 5 Median NDI pre- and post-treatment of groups A and B:

Outcome		Group A	Group B		
Measures		Median (IQR)	Median (IQR)	U- value	p-value
NDI (%)	Pretreatment	4 (4-2)	4 (4-4)	201	0.40
	Post-treatment	2 (4-2)	2 (4-2)	215.5	0.66
	Z-value	2.31	2.48		
		<i>p = 0.02*</i>	<i>p = 0.01*</i>		

IQR, Interquartile range; Z- value, Wilcoxon signed ranks test value; U- value, Mann-Whitney test value; p-value, Probability value. * indicate significance.


FIGURES

Figure 1: Flow diagram (CONSORT) of the progress through the study




Figure 2: A: Frog on the ground with upper limbs in adduction and open hip angle. B: Frog in the air with upper limbs in adduction and closed hip angle. C: Sitting with upper limbs in adduction and closed hip angle. D: Standing against the wall with upper limbs in adduction and open hip angle.




Figure 3: The measured angles from top to bottom: (1) Sagittal head tilt angle (gaze angle): the angle between a line between the tragus and the Canthus of the eye with the horizontal line, (2) Craniovertebral angle: angle formed between a line extending from C7 to the tragus of the ipsilateral ear and the horizontal line., and (3) Sagittal shoulder-C7 angle (rounded shoulder angle): the angle between a horizontal line through the lateral shoulder and another passing between C7 and the midpoint of the acromion (Singla et al., 2017). Interrupted lines represent the horizontal line.

Journal Pre-proof

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Journal Pre-proof